

Basic fuzzy logics and weak associative uninorms

E. Yang ¹

¹Department of Philosophy & Institute of Critical Thinking and Writing, Jeonbuk National University, Jeonju, Korea

eunsyang@jbnu.ac.kr

Abstract

Micanorm-based logics with a weak form of associativity are introduced and their completeness results are addressed. More concretely, first the basic wa_t -uninorm logic $\mathbf{WA}_t\mathbf{BUL}$ and its axiomatic extensions are introduced as $[0, t]$ -continuous wa_t -uninorm analogues of the logics based the $[0, 1]$ -continuous uninorms. Next algebraic structures characterizing the logics are introduced along with algebraic completeness results. Third, wa_t -uninorms are introduced as uninorms with weak t -associativity instead of associativity and associated properties are discussed. Finally, by virtue of Yang-style construction, it is verified that the logics based on wa_t -uninorms are complete on unit real interval $[0, 1]$, i.e., so called *standard* complete.

Keywords: Fuzzy logic, t-norm, wa_t -uninorm, uninorm, micanorm.

1 Introduction

The multiplication and addition of natural numbers are associative, whereas their division and subtraction are not. Associated with this, we may think of *partially* associative operations as operations between fully associative and fully non-associative operations. In fact, many such operations have been introduced. Especially, operations that satisfy weak forms of associativity have been addressed extensively. The following are most famous examples in mathematics: quasigroups ([?, ?]), hypergroups ([?]), power associativity in abstract algebra ([?, ?, ?]), hyperstructures ([?, ?, ?, ?, ?, ?]), weakly associative lattices ([?, ?, ?]) and weakly associative relation algebras ([?, ?, ?]).

In substructural fuzzy logic, logics based on more general structures have been introduced. T-norm-based logics ([?, ?, ?, ?]), uninorm-based logics ([?, ?, ?, ?, ?]), micanorm-based logics ([?, ?, ?, ?, ?, ?]) and mianorm-based logics ([?, ?, ?, ?, ?]) are examples. One interesting fact is that t-norm-based and uninorm-based logics require associativity as a structural rule, whereas micanorm-based and mianorm-based logics do not. Note that uninorms are micanorms with associativity. Then we may similarly consider binary operations satisfying weak forms of associativity as operations between uninorms and micanorms.

In fact, Yang [?, ?] introduced such operations as weak associative uninorms and dealt with corresponding logics. Especially he [?] introduced the fuzzy logics based on some particular weak t -associative uninorms (simply wa_t -uninorms), where the index ' t ' represents identity, $\mathbf{WA}_t\mathbf{BL}$, $\mathbf{WA}_t\mathbf{L}$, $\mathbf{WA}_t\mathbf{G}$, and $\mathbf{WA}_t\mathbf{\Pi}$ as a non-associative generalization of the continuous t-norm-based logics \mathbf{BL} (Basic fuzzy logic), \mathbf{L} (Łukasiewicz logic), \mathbf{G} (Gödel logic) and $\mathbf{\Pi}$ (Product logic), respectively, and showed that they are standard complete, i.e., complete over $[0, 1]$ (real unit interval). Those logics are $[0, t]$ -continuous wa_t -uninorm-based analogues of the continuous t-norm-based logics.

Notice that Gabbay and Metcalfe [?] introduced basic uninorm logics \mathbf{BUL} (Basic uninorm logic), \mathbf{IBUL} (Involutive BUL), \mathbf{CBUL} (Cancellative BUL) and \mathbf{CRL} (Cross ratio logic) as uninorm-based analogues of continuous t-norm-based logics. Those logics are based on $[0, 1]$ -continuous uninorms. Related to this fact, to provide a wa_t -uninorm generalization of the logics was given an open problem by Yang [?]. The problem can be divided into the following two questions.

- Q1: Can wa_t -uninorm analogues of the basic uninorm logics be introduced?

- Q2: Do those logics (if exist) form a $[0, t]$ -continuous wa_t -uninorm-based generalization of the $[0, 1]$ -continuous uninorm-based logics?

Notice also that a continuous t-norm \diamond has the properties

P1: if $o \leq l$, then $o = l \diamond (l \rightarrow o)$, for all $o, l \in [0, 1]$; and

P2: if $o \leq l \leq m$ and l is idempotent, then $o = o \diamond m$, for all $o, l, m \in [0, 1]$ (see [?]),

whereas a $[0, 1]$ -continuous uninorm \diamond has their restricted ones

P1': if $o \leq l < 1$, then $o = l \diamond (l \rightarrow o)$, for all $o, l \in [0, 1]$; and

P2': if u is $1 \rightarrow t$ and $o \leq u \leq l$, then $o = o \diamond l$, for all $o, l \in [0, 1]$ (see [?]).

We in general call the properties P1 and P1' *division* and *restricted division* properties, respectively. The element u in P2' plays the role of l in P2.

It would be interesting to mention that $[0, t]$ -continuous wa_t -uninorms introduced in [?] have the property P1' but need not have P2'. Note that as an idempotent element wa_t -uninorms need not require u . Then the following arises as a natural question.

- Q3: Does a wa_t -uninorm with the idempotent element u has the property P2' if u is added to a $[0, t]$ -continuous wa_t -uninorm?

As an answer to all of the three questions, we introduce basic wa_t -uninorm logics. More exactly, Sect. 2 introduces the basic wa_t -uninorm logic **WA_tBUL** and its axiomatic extensions as a $[0, t]$ -continuous wa_t -uninorm-based generalization, and addresses related algebraic semantics. Sect. 3 deals with some $[0, t]$ -continuous wa_t -uninorms as algebraic structures for the logics over $[0, 1]$. Sect. 4 provides standard completeness results for the logics by virtue of a construction of Yang-style in [?, ?].

We finally notice that a wa_t -uninorm has a subalgebra isomorphic to a t-norm. This implies at least two things. One is that since wa_t -uninorms have subalgebras isomorphic to t-norms, logically interesting t-norms such as Łukasiewicz, Gödel and Product t-norms and their applications to fuzzy logic and fuzzy sets can be still considered in wa_t -uninorms. The other is that since wa_t -uninorms are non-associative operations, properties of such operations and their applications such as compensation behavior and full reinforcement can be investigated in wa_t -uninorms.

2 Logics and algebraic semantics

Henceforth, the notion of *basic wa_t -uninorm-based logics* denotes substructural fuzzy logics being characterized by algebraic semantics, where the connective pair $(\&, \rightarrow)$ is interpreted by a basic wa_t -uninorm and its residuum. For these logics, this section consists of four steps: 1st. A language \mathcal{L} for the logics is provided; 2nd. On \mathcal{L} , the logics intended are introduced using a deductive consequence relation \vdash and related notions such as proof are considered; 3rd. Algebraic structures to characterize the logics are introduced and a related semantic consequence relation \models is defined along with related notions such as tautology and model; 4th. Using completeness theorem, it is verified that the deductive and semantic consequences are equivalent to each other.¹

We first introduce a language for basic wa_t -uninorm-based logics. A language \mathcal{L} for these logics is a countable propositional language with a set of formulas *For* built inductively by a set of variables *Var*, binary connectives \vee (disjunction), \wedge (conjunction), $\&$ (fusion), \rightarrow (implication), and constants $\top, \mathbf{t}, \mathbf{f}, \perp$. As the additional connectives and constant, it further has \leftrightarrow (biimplication), \sim (negation), \neg (negation) and U given by:

df1. $R \leftrightarrow P := (R \rightarrow P) \wedge (P \rightarrow R)$,

df2. $\sim R := R \rightarrow \mathbf{f}$,

df3. $\neg R := R \rightarrow \perp$, and

df4. $U := \top \rightarrow \mathbf{t}$.

We also simplify $R \wedge \mathbf{t}$ and $R \wedge U$ by $R_{\mathbf{t}}$ and R_U , respectively. An \mathcal{L} -*substitution* is a map $s : \text{For} \rightarrow \text{For}$ such that $s(\#(R_1, \dots, R_n)) = \#(s(R_1), \dots, s(R_n))$ for all n -ary connective $\#$ and all formulas $R_1, \dots, R_n \in \text{For}$. We denote the formulas by uppercase Latin letters R, P, Q, \dots , and their sets by uppercase Greek letters Γ, Σ, \dots

We next present axiomatizations for the logics using a deductive consequence relation \vdash .² (For the notion of logics in general as the same kind of objects as axiom systems, see [?].)

¹For substructural fuzzy logics and algebraic semantics in general, see [?, ?, ?].

²More exactly, the relation \vdash is introduced as a *consecution relation* since it is not necessary to be a proof or derivation relation from premises to conclusion in a logic. For this relation, see [?, ?].

Definition 2.1. 1. The basic weak t -associative uninorm logic $\mathbf{WA}_t\mathbf{BUL}$ consists of the following axioms and rules:

$R \rightarrow R$	(self-implication, SI)
$(R \wedge P) \rightarrow R, (R \wedge P) \rightarrow P$	(\wedge -elimination, \wedge -E)
$((R \rightarrow P) \wedge (R \rightarrow Q)) \rightarrow (R \rightarrow (P \wedge Q))$	(\wedge -introduction, \wedge -I)
$R \rightarrow (R \vee P), P \rightarrow (R \vee P)$	(\vee -introduction, \vee -I)
$((R \rightarrow Q) \wedge (P \rightarrow Q)) \rightarrow ((R \vee P) \rightarrow Q)$	(\vee -elimination, \vee -E)
$\perp \rightarrow R$	(ex falso quodlibet, EF)
$(R \& P) \rightarrow (P \& R)$	($\&$ -commutativity, $\&$ -C)
$(\mathbf{t} \rightarrow R) \leftrightarrow R$	(push and pop, PP)
$R \rightarrow (P \rightarrow (P \& R))$	($\&$ -adjunction, $\&$ -Adj)
$(R_t \& P_t) \rightarrow (R \wedge P)$	($\&$)
$(P \& (R \& (R \rightarrow (P \rightarrow Q)))) \rightarrow Q$	(residuation, Res')
$(R \rightarrow ((R \& (R \rightarrow P)) \& (P \rightarrow Q))) \rightarrow (R \rightarrow Q)$	(transitivity, T')
$((Z \& W) \rightarrow (Z \& (W \& (R \rightarrow P)_t))) \vee (Z' \rightarrow (W' \rightarrow ((W' \& Z') \& (P \rightarrow R)_t)))$	(prelinearity, PL)
$(R_t \& (P_t \& Q_t)) \leftrightarrow ((R_t \& P_t) \& Q_t)$	(weak \mathbf{t} -associativity, wAS_t)
$(\mathbf{t} \rightarrow R) \vee (R \rightarrow P) \vee (P \rightarrow (R \& (R \rightarrow P)))$	(weak \mathbf{t} -restricted divisibility, $RDIV_t^w$)
$R_U \rightarrow (U \& R_U)$	(U -restricted identity, U -RI)
$(U \vee R_t) \rightarrow ((U \vee R_t) \& (U \vee R_t))$	(\mathbf{t}_U -idempotence, ID_U^t)
$R \rightarrow P, R \vdash P$	(modus ponens, mp)
$R \vdash R_t$	(\mathbf{t} -adjunction, adj_t)
$R \vdash (Z \& W) \rightarrow (Z \& (W \& R))$	($\&$ -associativity, $\&$ -as)
$R \vdash Z \rightarrow (W \rightarrow ((W \& Z) \& R))$	(\rightarrow -associativity, \rightarrow -as)

2. The following are logics extending $\mathbf{WA}_t\mathbf{BUL}$:

- $\mathbf{WA}_t\mathbf{IBUL}$ (Involutive $\mathbf{WA}_t\mathbf{BUL}$) is $\mathbf{WA}_t\mathbf{BUL}$ plus $\sim \sim R \rightarrow R$ (double negation elimination, DNE); and $(\neg \neg R)_U \rightarrow R$ (U -double negation elimination, DNE_U).
- $\mathbf{WA}_t\mathbf{CBUL}$ (Cancellative $\mathbf{WA}_t\mathbf{BUL}$) is $\mathbf{WA}_t\mathbf{BUL}$ plus $(R \wedge \neg R) \rightarrow \perp$ (Gödel negation, GN); $(R \rightarrow P) \rightarrow (\neg P \rightarrow \neg R)$ (contraposition, CP); and $(\neg \neg R)_U \rightarrow ((R_U \rightarrow (R_U \& P))_U \rightarrow P)$ (weak U -restricted cancellation, $RCAN_U^w$).
- $\mathbf{WA}_t\mathbf{GBUL}$ (Restricted idempotent $\mathbf{WA}_t\mathbf{BUL}$) is $\mathbf{WA}_t\mathbf{BUL}$ plus $(R \& R)_U \leftrightarrow R_U$ (U -restricted idempotence, RID_U).
- $\mathbf{WA}_t\mathbf{CRL}$ (Cross ratio $\mathbf{WA}_t\mathbf{BUL}$) is $\mathbf{WA}_t\mathbf{IBUL}$ plus $\mathbf{t} \leftrightarrow \mathbf{f}$ (fixed-point, FP).

Remark 2.2. 1. We obtain the $[0, t]$ -continuous $w\mathbf{a}_t$ -uninorm logic $\mathbf{WA}_t\mathbf{BL}$ by dropping (U -RI) and (ID_U^t) from $\mathbf{WA}_t\mathbf{BUL}$, the $w\mathbf{a}_t$ -uninorm logic $\mathbf{WA}_t\mathbf{MUL}$ by eliminating ($RDIV_t^w$) from $\mathbf{WA}_t\mathbf{BL}$, and the mininorm logic \mathbf{MICAL} , described as SL_e^ℓ in [?, ?, ?], by dropping (wAS_t) from $\mathbf{WA}_t\mathbf{MUL}$ (see [?, ?, ?]).

2. The systems $\mathbf{WA}_t\mathbf{BUL}$, $\mathbf{WA}_t\mathbf{IBUL}$, $\mathbf{WA}_t\mathbf{CBUL}$, and $\mathbf{WA}_t\mathbf{CRL}$ are $[0, t]$ -continuous $w\mathbf{a}_t$ -uninorm analogues of the $[0, 1)$ -continuous uninorm logics \mathbf{BUL} , \mathbf{IBUL} , \mathbf{CBUL} , and \mathbf{CRL} , respectively, which are obtained as follows (see [?]):

- Basic uninorm logic \mathbf{BUL} is \mathbf{UL} plus $df1$ to $df4$ and: $U \leftrightarrow (U \& U)$ (U -idempotence, U -ID).
 $(\top \rightarrow R) \vee (R \rightarrow (P \wedge U)) \vee (P \rightarrow (R \& (R \rightarrow P)))$ (restricted divisibility, $RDIV$).
- \mathbf{IBUL} (Involutive \mathbf{BUL}) is \mathbf{BUL} plus (DNE).
- \mathbf{CBUL} (Cancellative \mathbf{BUL}) is \mathbf{BUL} plus $(R \rightarrow \perp) \vee (\top \rightarrow R) \vee ((R \rightarrow (R \& P)) \rightarrow P)$ (restricted cancellation, $RCAN$).
- \mathbf{CRL} (Cross ratio logic) is \mathbf{IBUL} plus (FP).

Definition 2.3. $\mathbf{WA}_t\mathbf{Ls} = \{\mathbf{WA}_t\mathbf{BUL}, \mathbf{WA}_t\mathbf{IBUL}, \mathbf{WA}_t\mathbf{CBUL}, \mathbf{WA}_t\mathbf{GBUL}, \mathbf{WA}_t\mathbf{CRL}\}$.

We define a proof as a deductive consequence relation \vdash , which is a relation between sets of formulas and formulas. A *proof* in a theory Σ (a set of formulas) on $\mathbf{WA}_t\mathbf{L}$ ($\in \mathbf{WA}_t\mathbf{Ls}$) is defined as a sequence of formulas whose elements are either members of Σ , axioms of $\mathbf{WA}_t\mathbf{L}$, or are derived from previous elements of the sequence by rules of $\mathbf{WA}_t\mathbf{L}$. By $\Sigma \vdash_{\mathbf{WA}_t\mathbf{L}} R$, we mean that R is *provable* in Σ on $\mathbf{WA}_t\mathbf{L}$.

A p -formula is constructed by $Var \cup \{p\}$ and a p -substitution is a substitution in the extended language. Let R be a formula, Z be a p -formula and s be a p -substitution such that $s(p) = R$ and $s(q) = q$ for $q \in Var$. Let $Z(R)$ denote the formula $s(Z)$, which is in the original set of variables. For a set Σ of p -formulas, Σ^* is defined as the least set of p -formulas, where $p \in \Sigma^*$ and $Z(P) \in \Sigma^*$ for all $P \in \Sigma^*$ and all $Z \in \Sigma$, and $\Pi(\Sigma)$ is the least set of p -formulas containing $\Sigma \cup \{\mathbf{t}\}$ and closed under $\&$. Let bDT , the set of basic deduction terms, be a set of p -formulas. A substructural logic L is almost mp -based on bDT if (i) the set bDT is closed under every p -substitution s , $s(p) = p$, (ii) L is an axiomatic system with the rules (mp) and those from $\{\langle R, P(R) \rangle : R \in For \text{ of } L \text{ and } P \in bDT\}$, and (iii) for all $Q \in bDT$ and all formulas R and P , one has some $Q_1, Q_2 \in bDT^*$ such that $\vdash_L Q_1(R \rightarrow P) \rightarrow (Q_2(R) \rightarrow Q(P))$ (see ([?, ?, ?, ?])).

Since WA_tL is an almost mp -based logic, we have the following deduction theorem for WA_tL .

Theorem 2.4. [?, ?, ?] *Given a set $\Gamma \cup \{R, P\}$ of formulas, it holds:*

$$\Sigma, R \vdash_{WA_tL} P \text{ if and only if } \Gamma \vdash_{WA_tL} Q(R) \rightarrow P \text{ for some } Q \in \Pi(bDT^*).$$

We then introduce algebraic structures corresponding to WA_tL and define a semantic consequence relation \models . For convenience, the notations ' \top ', ' \perp ' are ambiguously used as both constants and special elements, and similarly the notations ' \neg ', ' \sim ', ' \rightarrow ', ' \vee ', and ' \wedge ' as both connectives and operators.

Definition 2.5. 1. [?] *A bounded residuated lattice-ordered pointed commutative groupoid with identity is a structure $(S, \perp, \top, f, t, \rightarrow, \star, \vee, \wedge)$ such that:*

- $(S, \perp, \top, \vee, \wedge)$ is a bounded lattice, where the least and greatest elements are \perp and \top , respectively.
- (S, \star, t) is a commutative groupoid with identity.
- f is a special element in S .
- $l \leq o \rightarrow m$ if and only if $o \star l \leq m$, for all $o, l, m \in S$ (residuation).

2. [?] *Let o_t be $o \wedge t$. A residuated wa_t -monoidal lattice is a bounded residuated lattice-ordered pointed commutative groupoid with identity that satisfies:*

- $o_t \star (l_t \star m_t) = (o_t \star l_t) \star m_t$, for all $o, l, m \in S$ (wa_t^S).

3. [?] *A WA_tMUL -algebra is a residuated wa_t -monoidal lattice satisfying:*

- $t \leq ((m \star p) \rightarrow (m \star (p \star (o \rightarrow l)_t))) \vee (m' \rightarrow (p' \rightarrow ((p' \star m') \star (l \rightarrow o)_t)))$, for all $o, l, m, p, m', p' \in A$ (PL^S).

Definition 2.6. (WA_tL -algebras) *Let $\sim o$, $\neg o$, and u be $o \rightarrow f$, $o \rightarrow \perp$ and $\top \rightarrow t$, respectively.*

1. *A WA_tBUL -algebra is a WA_tMUL -algebra satisfying:*

- $t \leq (t \rightarrow o) \vee (o \rightarrow l) \vee (l \rightarrow (o \star (o \rightarrow l)))$, for all $o, l \in S$ ($RDIV_t^{wS}$),
- $o_u \leq (u \star o_u)$, for all $o \in S$ ($U-RF^S$), and
- $(u \vee o_t) \leq ((u \vee o_t) \star (u \vee o_t))$, for all $o \in S$ (ID_U^tS).

2. *A WA_tIBUL -algebra is a WA_tBUL -algebra satisfying:*

- $t \leq \sim \sim o \rightarrow o$, for all $o \in S$ (DNE^S), and
- $t \leq (\neg \neg o)_u \rightarrow o$, for all $o \in S$ (DNE_U^S)

3. *A WA_tCBUL -algebra is a WA_tBUL -algebra satisfying:*

- $t \leq (o \wedge \neg o) \rightarrow \perp$ for all $o \in S$ (GN^S)
- $t \leq (o \rightarrow l) \rightarrow (\neg l \rightarrow \neg o)$ for all $o, l \in S$ (CP^S)
- $t \leq (\neg \neg o)_u \rightarrow ((o_u \rightarrow (o_u \star l))_u \rightarrow l)$ for all $o, l \in S$ ($RCAN_U^{wS}$).

4. *A WA_tGBUL -algebra is a WA_tBUL -algebra satisfying*

- $(o \star o)_u = o_u$, for all $o \in S$ (RID_U^S).

5. *A WA_tCRL -algebra is a WA_tIBUL -algebra satisfying*

- $t = f$ (FP^S).

All of these algebras are called WA_tL -algebras.

Let \mathcal{S} be a WA_tL -algebra, \mathcal{K} be a class of WA_tL -algebras, R and P be formulas and Γ be a theory. We then define several notions for the semantic consequence relation for \mathbf{WA}_tL . An \mathcal{S} -interpretation is a homomorphism $It : For \rightarrow \mathcal{S}$ such that $It(\perp) = \perp$, $It(\top) = \top$, $It(\mathbf{t}) = t$, $It(\mathbf{f}) = f$, $It(R \rightarrow P) = It(R) \rightarrow It(P)$, $It(R \& P) = It(R) \star It(P)$, $It(R \vee P) = It(R) \vee It(P)$, $It(R \wedge P) = It(R) \wedge It(P)$ (and so $It(\sim R) = \sim It(R)$, $It(\neg R) = \neg It(R)$ and $It(U) = u$). R is an t -tautology in \mathcal{S} , an \mathcal{S} -tautology for simplicity, if $t \leq It(R)$ for every \mathcal{S} -interpretation It . An \mathcal{S} -interpretation It is an \mathcal{S} -model of Γ if $t \leq It(R)$ for all $R \in \Gamma$. The class of \mathcal{S} -models of Γ is referred to by $Mod(\Gamma, \mathcal{S})$. R is a semantic consequence of Γ over \mathcal{K} , $\Gamma \models_{\mathcal{K}} R$, if $Mod(\Gamma, \mathcal{S}) = Mod(\Gamma \cup \{R\}, \mathcal{S})$ for each $\mathcal{S} \in \mathcal{K}$. \mathcal{S} is a \mathbf{WA}_tL -algebra if and only if $\Gamma \vdash_{WA_tL} R$, $WA_tL \in WA_tLs$, entails that $\Gamma \models_{\{\mathcal{S}\}} R$, \mathcal{S} a corresponding WA_tL -algebra. $MOD(WA_tL)$ and $MOD^l(WA_tL)$ refer to the class of \mathbf{WA}_tL -algebras and the class of linearly ordered \mathbf{WA}_tL -algebras, respectively. For simplicity, $\Gamma \vdash_{WA_tL} R$ and $\Gamma \models_{WA_tL}^l R$ express $\Gamma \models_{MOD(WA_tL)} R$ and $\Gamma \models_{MOD^l(WA_tL)} R$, respectively.

We finally prove completeness for WA_tL , which shows that the two consequence relations \vdash and \models on WA_tL are equivalent to each other.

Theorem 2.7. 1. (Strong completeness) Suppose that Γ is a theory over $WA_tL \in \{\mathbf{WA}_tBUL, \mathbf{WA}_tGBUL\}$ and R be a formula. $\Gamma \vdash_{WA_tL} R$ if and only if $\Gamma \models_{WA_tL} R$ if and only if $\Gamma \models_{WA_tL}^l R$.

2. (Finite strong completeness) Let Γ be a finite theory over $WA_tL \in WA_tLs$ and R be a formula. $\Gamma \vdash_{WA_tL} R$ if and only if $\Gamma \models_{WA_tL} R$ if and only if $\Gamma \models_{WA_tL}^l R$.

Proof. The claims are directly obtained by Theorem 3.1.8 in [?]. □

3 Basic weak- t -associative uninorms

We henceforth express \perp and \top on $[0, 1]$ by 0 and 1, respectively. As usual, we call a WA_tL -algebra on $[0, 1]$ standard. This section consists of two steps: 1st. Basic notions of weak- t -associative uninorms (wa_t -uninorms for simplicity) as standard WA_tL -algebras are introduced. 2nd. It is verified that such wa_t -uninorms form WA_tL -algebra on $[0, 1]$.

The operator \diamond is a micanorm in standard WA_tL -algebras. A micanorm is a function $\diamond : [0, 1]^2 \rightarrow [0, 1]$ satisfying: for all $o, l, m \in [0, 1]$ and for some $t \in [0, 1]$,

- (i) $o \diamond l = l \diamond o$ (commutativity),
- (ii) $t \diamond o = o = o \diamond t$ (identity), and
- (iii) $o \leq l$ entails $o \diamond m \leq l \diamond m$ and $m \diamond o \leq m \diamond l$ (isotonicity), see [?].

A weak- t -associative uninorm (wa_t -uninorm for simplicity) is a micanorm satisfying:

$$(wAS_t) \ o, l, m \leq t \text{ entails } o \diamond (l \diamond m) = (o \diamond l) \diamond m, \text{ see [?].}$$

Note that a uninorm is a micanorm satisfying: for all $o, l, m \in [0, 1]$,

$$(AS) \ o \diamond (l \diamond m) = (o \diamond l) \diamond m,$$

and a t -norm is a uninorm satisfying $t = 1$. We call a micanorm \diamond satisfying $0 \diamond 1 = 1 \diamond 0 = 0$ conjunctive, and a micanorm \diamond satisfying $0 \diamond 1 = 1 \diamond 0 = 1$ disjunctive (see [?]).

Note also that we can get the implication functions using residual operators of micanorms and wa_t -uninorms. We call a micanorm \diamond residuated whenever we have a function $\rightarrow : [0, 1]^2 \rightarrow [0, 1]$ such that it satisfies (residuation), called the residuum of \diamond . Given a micanorm (wa_t -uninorm resp) \diamond , implication \rightarrow can be defined as follows: for all $o, l \in [0, 1]$,

$$o \rightarrow l := \sup\{m \in [0, 1] : o \diamond m \leq l\}.$$

Over $[0, 1]$, the operation \diamond of a UL-algebra is a conjunctive uninorm with its residuum \rightarrow and identity t , and conversely a residuated uninorm \diamond arises a UL-algebra (see [?]). Similarly for a MICAL-algebra on $[0, 1]$ and a residuated micanorm (see [?]).

Let the wa_t -uninorms to characterize WA_tL -algebras be basic wa_t -uninorms. We notice that an arbitrary residuated $[0, 1]$ -continuous uninorm \diamond arises a BUL-algebra on $[0, 1]$ (see [?]). Analogously, some residuated $[0, t]$ -continuous wa_t -uninorm \diamond arises a WA_tL -algebra on $[0, 1]$. We verify it.

A negation function $\nu : [0, 1] \rightarrow [0, 1]$ is in general defined as a non-increasing function which satisfies $\nu(1) = 0$ and $\nu(0) = 1$. Since this definition does not work for uninorms in general³, we instead define a non-increasing function as a negation. A negation ν is called: conjunctive if it satisfies:

³For instance, a non-increasing function ν needs not satisfy $\nu(1) = 0$ in a uninorm.

- (conj) $\nu(0) = 1$;

fixed-pointed if it satisfies:

- (fp) $\nu(id) = id$;

super-involutive if it satisfies:

- (s-inv) $\nu^2(o) \geq o$ for all $o \in [0, 1]$;

weak if it is super-involutive and satisfies:

- (w) $\nu(0) = 1$ and $\nu(1) = 0$;

involutive if it is weak and satisfies:

- (inv) $\nu^2(o) = o$ for all $o \in [0, 1]$.

It is known that involutive negations on $[0, 1]$ are isomorphic to each other (see [?]). Thus for all $o \in [0, 1]$, an involutive negation $\nu(o)$ is definable by $1 - o$. The negation defined by $1 - o$ is called *standard*. An involutive negation ν on $[0, u]$ is called *u-involutive*.

WA_tL -algebras on $[0, 1]$ can be obtained from residuated $[0, t]$ -continuous wa_t -uninorms satisfying:

- (t_u -ID) $u \leq o \leq t$ entails $o = (o \diamond o)$, for all $o \in [0, 1]$, and
- (u -RI) $o \leq u$ entails $o = (u \diamond o)$, for all $o \in [0, 1]$.

We call wa_t -uninorms satisfying these conditions t_u -idempotent restricted u -identical wa_t -uninorms. These wa_t -uninorms form basic wa_t -uninorms.

Proposition 3.1. *Let u be $1 \rightarrow t$.*

1. A structure $([0, 1], 0, 1, t, f, u, \diamond, \max, \min, \rightarrow)$ is a standard WA_tBUL -algebra if \diamond is a residuated $[0, t]$ -continuous t_u -idempotent restricted u -identical wa_t -uninorm with identity t .
2. A structure $([0, 1], 0, 1, t, f, u, \nu_1, \nu_2, \diamond, \max, \min, \rightarrow)$ is a standard WA_tIBUL -algebra if \diamond is a residuated $[0, t]$ -continuous t_u -idempotent restricted u -identical wa_t -uninorm with identity t , an involutive negation ν_1 and a u -involutive negation ν_2 .
3. Suppose that ν is a negation satisfying (GN^S) , i.e., Gödel negation. A structure $([0, 1], 0, 1, t, f, u, \nu, \diamond, \max, \min, \rightarrow)$ is a standard WA_tCBUL -algebra if \diamond is a residuated $[0, t]$ -continuous t_u -idempotent restricted u -identical wa_t -uninorm with identity t and Gödel negation ν , and is strictly isotone on $[0, u]$.
4. A structure $([0, 1], 0, 1, t, f, u, \diamond, \max, \min, \rightarrow)$ is a standard WA_tGBUL -algebra if \diamond is a residuated $[0, t]$ -continuous $[0, t]$ -idempotent wa_t -uninorm with identity t .
5. A structure $([0, 1], 0, 1, \frac{1}{2}, \frac{1}{2}, u, \nu_1, \nu_2, \diamond, \max, \min, \rightarrow)$ is a standard WA_tCRL -algebra if \diamond is a residuated $[0, t]$ -continuous t_u -idempotent restricted u -identical wa_t -uninorm with a standard negation ν_1 , a u -involutive negation ν_2 and identity $t = \frac{1}{2} = \nu_1(t)$.

Proof. 1. Let \diamond be a residuated $[0, t]$ -continuous t_u -idempotent restricted u -identical wa_t -uninorm with t as identity. Since one is capable of obtaining a WA_tBL -algebra from an arbitrary residuated $[0, t]$ -continuous wa_t -uninorm (see [?]), we just need to verify that $(U-RI^S)$ and $(ID_U^t)^S$ hold. For $(U-RI^S)$, one has to show that: for all $o \in [0, 1]$,

$$\min\{o, u\} \leq u \diamond \min\{o, u\}.$$

Let $o \leq u$. Since \diamond is restricted u -identical, we obtain $o \leq u \diamond o$ by $(u-RI)$. Otherwise, $\min\{o, u\} = u$ and so $u \leq u \diamond u$ by $(u-RI)$. Hence the claim holds.

For $(ID_U^t)^S$, one has to show that: for all $o \in [0, 1]$,

$$\max\{u, \min\{o, t\}\} \leq \max\{u, \min\{o, t\}\} \diamond \max\{u, \min\{o, t\}\}.$$

Let $u \leq o \leq t$. Then $\max\{u, \min\{o, t\}\} = o$. Also, since \diamond is t_u -idempotent, we obtain $o = o \diamond o$ and so $o \leq o \diamond o$. Let $t < o$. Then $\max\{u, \min\{o, t\}\} = t$ and so $t \leq t \diamond t$. Otherwise, $\max\{u, \min\{o, t\}\} = u$. Then, since \diamond is t_u -idempotent, we obtain $u = u \diamond u$ and so $u \leq u \diamond u$. Hence the claim holds.

2. Let \diamond be a residuated $[0, t]$ -continuous t_u -idempotent restricted u -identical wa_t -uninorm with identity t , an involutive negation ν_1 and a u -involutive negation ν_2 . We need to further verify that (DNE^S) and (DNE_U^S) hold. For (DNE^S) , one has to show that: for all $o \in [0, 1]$,

$$t \leq \nu_1^2(o) \rightarrow o.$$

Since ν_1 is involutive, i.e., $\nu_1^2(o) = o$, we have $\nu_1^2(o) \leq o$ and so $t \leq \nu_1^2(o) \rightarrow o$ by (residuation).

For $(\text{DNE}_U^{\mathcal{S}})$, one has to show that: for all $o \in [0, 1]$,

$$t \leq \min\{\nu_2^2(o), u\} \rightarrow o.$$

Let $o \leq u$. Since ν_2 is u -involutive, we have $\min\{\nu_2^2(o), u\} = \nu_2^2(o) = o$. Then we further obtain $\nu_2^2(o) \leq o$ and so $t \leq \nu_2^2(o) \rightarrow o$ by (residuation). Otherwise, i.e., if $u < o$, then $u = u \diamond t < o$ and so $t \leq u \rightarrow o$ by (residuation). Moreover we have $\min\{\nu_2^2(o), u\} = u$. Hence the claim holds.

3. Let \diamond be a residuated $[0, t]$ -continuous t_u -idempotent restricted u -identical wa_t -uninorm with identity t and Gödel negation ν , and is strictly isotone on $[0, u]$. We need to further verify that $(\text{GN}^{\mathcal{S}})$, $(\text{CP}^{\mathcal{S}})$ and $(\text{RCAN}_U^{\mathcal{S}})$ hold.

For $(\text{GN}^{\mathcal{S}})$, one has to show that: for all $o \in [0, 1]$,

$$t \leq \min\{o, \nu(o)\} \rightarrow o.$$

By (residuation), we may instead show that $\min\{o, \nu(o)\} \leq o$. If $o = 0$, then clearly $\min\{o, \nu(o)\} \leq o$. Otherwise, $\nu(o) = 0$ and so $\min\{o, \nu(o)\} \leq o$.

For $(\text{CP}^{\mathcal{S}})$, one has to show that: for all $o, l \in [0, 1]$,

$$t \leq (o \rightarrow l) \rightarrow (\nu(l) \rightarrow \nu(o)).$$

Since the implication \rightarrow is transitive, it holds that

$$t \leq (o \rightarrow l) \rightarrow ((l \rightarrow 0) \rightarrow (o \rightarrow 0)).$$

Hence the claim holds by df3.

For $(\text{RCAN}_U^w{}^{\mathcal{S}})$, one has to show that: for all $o, l \in [0, 1]$,

$$t \leq \min\{\neg\neg o, t\} \rightarrow (\min\{(\min\{o, u\} \rightarrow (\min\{o, u\} \diamond l)), u\} \rightarrow l).$$

Let $o = 0$. We have $\neg\neg o = 0$. Thus the claim holds. Let $o \neq 0$. We obtain $\neg\neg o = 1$ and so $\min\{\neg\neg o, t\} = t$. Then, by (residuation) (twice), it suffices to verify

$$(\alpha) \min\{\min\{o, u\} \rightarrow (\min\{o, u\} \diamond l), u\} \leq l.$$

Here, we consider the case $o \leq u$. If $u < l$, then (α) holds because $\min\{o \rightarrow (o \diamond l), u\} \leq l$. Let $l < u$. Then, since $o \diamond l \leq \min\{o, l\} \leq l$ and \diamond is strict isotone on $[0, u]$, (α) also holds.

4. Let \diamond be a residuated $[0, t]$ -continuous $[0, t]$ -idempotent wa_t -uninorm with identity t . We need to further verify that $(U\text{-RI}^{\mathcal{S}})$, $(\text{ID}_U^t{}^{\mathcal{S}})$ and $(\text{RID}_U^{\mathcal{S}})$ hold. For $(U\text{-RI}^{\mathcal{S}})$, one may verify $(u\text{-RI})$. Let $o \leq u$. Then since \diamond is $[0, t]$ -idempotent, we have

$$o = o \diamond o \leq u \diamond o \leq t \diamond o = o,$$

and so $o = (u \diamond o)$. Hence $(u\text{-RI})$ holds.

For $(\text{ID}_U^t{}^{\mathcal{S}})$, one may verify $({}^t_u\text{-ID})$. Let $u \leq o \leq t$. Then since \diamond is $[0, t]$ -idempotent, we have $o = o \diamond o$. Hence $({}^t_u\text{-ID})$ holds.

For $(\text{RID}_U^{\mathcal{S}})$, one has to show that: for all $o \in [0, 1]$,

$$(o \diamond o)_u = o_u.$$

If $u \leq o$, then $(o \diamond o)_u = u = o_u$. Otherwise, $(o \diamond o)_u = o = o_u$ since \diamond is $[0, t]$ -idempotent. Hence the claim holds.

5. Let \diamond be a residuated $[0, t]$ -continuous t_u -idempotent restricted u -identical wa_t -uninorm with a standard negation ν_1 , a u -involutive negation ν_2 and identity $t = \frac{1}{2} = \nu_1(t)$. Since $t = \frac{1}{2} = \nu_1(t) = f$, the claim directly follows from 2. \square

4 Standard completeness

Standard completeness results are established for $\mathbf{WA}_t\mathbf{L}$ using a construction of Yang–style in [?, ?]. The idea for standard completeness is this: Countable or finite linearly ordered $\mathbf{WA}_t\mathbf{L}$ -algebras are embeddable into dense linearly ordered ones, and these algebras again embeddable into standard algebras. This section consists of four steps: 1st. Standard completeness of $\mathbf{WA}_t\mathbf{BUL}$ is provided based on the above idea. 2nd. This proof is applied to its extensions. 3rd. It is verified that $\mathbf{WA}_t\mathbf{L}$ -algebras on $[0, 1]$ have subalgebras isomorphic to corresponding t -norms. 4th. It is verified that standard $\mathbf{WA}_t\mathbf{L}$ -algebras are given on restricted v -identical ι_v -idempotent $[0, t]$ -continuous wa_t -uninorms.

Notice first that linearly ordered countable or finite \mathbf{MICAL} -algebras can be embedded into a standard algebra. (For the shake of easiness, we add the ‘less than or equal to’ relation symbol “ \leq .”)

Fact 4.1. ([?]) *For each linearly ordered countable or finite \mathbf{MICAL} -algebra $\mathbf{S} = (S, \leq_S, \perp, \top, f, t, \rightarrow, \star, \vee, \wedge)$, one can construct a countable ordered set T , a binary operation \ominus on T , and a function i from S into T satisfying:*

1. T is a dense ordered set and has a minimum \wedge , a maximum \vee and special elements ρ and ι .
2. $(T, \ominus, \lesssim, \iota)$ is a commutative, isotone linearly ordered groupoid with identity.
3. \ominus is left-continuous and conjunctive over the order topology on (T, \lesssim) .
4. i is an embedding function of the structure $(S, \leq_S, \perp, \top, f, t, \rightarrow, \star, \vee, \wedge)$ into $(T, \lesssim, \wedge, \vee, \rho, \iota, \ominus, \max, \min)$, and for all $o, l \in S$, $i(o \rightarrow l)$ is the residuum of $i(o)$ and $i(l)$ in $(T, \lesssim, \wedge, \vee, \rho, \iota, \ominus, \max, \min)$.

We first establish standard completeness for $\mathbf{WA}_t\mathbf{BUL}$.

Proposition 4.2. *For each countable or finite, linearly ordered $\mathbf{WA}_t\mathbf{BUL}$ -algebra $\mathbf{S} = (S, \leq_S, \perp, \top, f, t, u, \rightarrow, \star, \vee, \wedge)$, one can construct a countable ordered set T , a binary operation \ominus on T , and a function i from S into T satisfying:*

1. T is densely ordered and has a minimum \wedge , a maximum \vee , and special elements ι , ρ and v .
2. $(T, \ominus, \lesssim, \iota, v)$ is a linearly ordered, isotonic, commutative ι_v -idempotent restricted v -identical weak ι -associative groupoid with identity ι .
3. \ominus is left-continuous and conjunctive over the order topology on (T, \lesssim) , and continuous over $\{o \in T : \wedge \lesssim o \lesssim \iota\}$.
4. i is an embedding function of the structure $(S, \leq_S, \top, \perp, f, t, u, \wedge, \vee, \star)$ into $(T, \lesssim, \vee, \wedge, \rho, \iota, v, \min, \max, \ominus)$, and for all $\sigma, \theta \in S$, $i(\sigma \rightarrow \theta)$ is the residuum of $i(\sigma)$ and $i(\theta)$ in $(T, \lesssim, \vee, \wedge, \rho, \iota, v, \min, \max, \ominus)$.

Proof. For easiness, let S be a subset of $[0, 1] \cap \mathbf{Q}$ with countable or finite elements, where 0 and 1 are the least and greatest elements, each of which corresponds to \perp and \top , respectively. Let

$$T = \{(\sigma, o) : \sigma \in S \setminus \{0 (= \perp)\} \text{ and } o \in \mathbf{Q} \cap (0, \sigma]\} \cup \{(0, 0)\}.$$

For $(\sigma, o), (\theta, l) \in T$, we define:

$$(\sigma, o) \lesssim (\theta, l) \text{ if and only if either } \sigma <_S \theta, \text{ or } \sigma =_S \theta \text{ and } o \leq l.$$

Let ρ , ι and v be (f, f) , (t, t) and (u, u) , respectively. Since $\mathbf{WA}_t\mathbf{BUL}$ -algebras are \mathbf{MICAL} -algebras, we just need to verify that $(T, \ominus, \lesssim, \iota, v)$ is a ι_v -idempotent restricted v -identical weak ι -associative groupoid with identity in 2 and \ominus is continuous over $\{o \in T : \wedge \lesssim o \lesssim \iota\}$ in 3. For convenience, the index S in \leq_S and $=_S$ is dropped unless specified otherwise.

Define, for $(\sigma, o), (\theta, l) \in T$:

$$(\sigma, o) \ominus (\theta, l) = \begin{cases} \max\{(\sigma, o), (\theta, l)\} & \text{if } \sigma \star \theta = \sigma \vee \theta, \sigma \neq \theta, \text{ and} \\ & (\sigma, o) \lesssim \iota \text{ or } (\theta, l) \lesssim \iota; \\ \min\{(\sigma, o), (\theta, l)\} & \text{if } \sigma \star \theta = \sigma \wedge \theta, \text{ and} \\ & (\sigma, o) \lesssim \iota \text{ or } (\theta, l) \lesssim \iota; \\ (\sigma \star \theta, \sigma \star \theta) & \text{otherwise.} \end{cases}$$

Let $T|_\iota := \{(\sigma, o) \in T : (0, 0) \lesssim (\sigma, o) \lesssim (t, t)\}$. For weak ι -associativity and continuity on $T|_\iota$ of \ominus , see Proposition 3 in [?]. For restricted v -identity, we have to show (u -RI), i.e.,

$$\text{if } (\sigma, o) \lesssim v, \text{ then } (\sigma, o) = v \ominus (\sigma, o).$$

Let $(\sigma, o) \lesssim v = (u, u)$. Then $\sigma \leq u$ and so $\sigma = u \star \sigma$. Thus we have $(\sigma, o) = \min\{(u, u), (\sigma, o)\} = v \ominus (\sigma, o)$. Hence the claim holds.

For ι_v -idempotence, we have to show $(\iota_v\text{-ID})$, i.e.,

$$\text{if } v \lesssim (\sigma, o) \lesssim \iota, \text{ then } (\sigma, o) = (\sigma, o) \ominus (\sigma, o).$$

Let $v \lesssim (\sigma, o) \lesssim \iota$. Then $u \leq \sigma \leq t$ and so $\sigma = \sigma \star \sigma$. Thus we have $(\sigma, o) = \min\{(\sigma, o), (\sigma, o)\} = (\sigma, o) \ominus (\sigma, o)$. Hence the claim holds.

The remaining conditions can be proved as in Proposition 2 in [?]. \square

Proposition 4.3. *Each dense countable linearly ordered $\mathbf{WA}_t\mathbf{BUL}$ -algebra is embeddable into a standard $\mathbf{WA}_t\mathbf{BUL}$ -algebra.*

Proof. Let first T, \mathbf{S} , etc. be given as in Proposition ???. Since (T, \lesssim) is a dense linearly ordered countable set with least and greatest elements, (T, \lesssim) is order isomorphic to $([0, 1] \cap \mathbf{Q}, \leq)$. Also, let f be such an ordered isomorphism, and for $o, l \in [0, 1]$, $o \ominus' l = f(f^{-1}(o) \ominus f^{-1}(l))$ and, for all $o \in S$, $i'(o) = f(i(o))$. Then if the conditions 1, 2, 3, and 4 hold, then the structure $([0, 1] \cap \mathbf{Q}, \leq, 0, 1, \rho, \iota, v, \ominus')$ and i' satisfy conditions 1, 2, 3, and 4 of Proposition ??? whenever the structure $(T, \lesssim, \wedge, \vee, \rho, \iota, v, \ominus)$ and i do the same. Thus we may assume $T = [0, 1] \cap \mathbf{Q}$, $\lesssim = \leq$ and $\ominus = \ominus'$.

Define: for all $o, l \in [0, 1]$,

$$o \hat{\ominus} l = \sup_{v \in T: v \leq o} \sup_{w \in T: w \leq l} v \ominus w.$$

Since $\mathbf{WA}_t\mathbf{BUL}$ -algebras are \mathbf{MICAL} -algebras, we just need to verify that $\hat{\ominus}$ is restricted v -identical, ι_v -idempotent, weak ι -associative, and continuous over $[0, \iota]$. For weak ι -associativity and continuity over $[0, \iota]$ of $\hat{\ominus}$, see Proposition 4 in [?]. We prove restricted v -identity and ι_v -idempotence of $\hat{\ominus}$.

For restricted v -identity of $\hat{\ominus}$, we need to verify that: for all $o \in T$,

$$\text{if } o \lesssim v, \text{ then } o = v \hat{\ominus} o.$$

Let $o \lesssim v$. Then we have $o = v \ominus o$ and so

$$\sup_{r \in T: r \leq o} r = \sup_{s \in T: s \leq v} \sup_{r \in T: r \leq o} s \ominus r.$$

Hence the claim holds.

For ι_v -idempotence of $\hat{\ominus}$, we need to verify that: for all $o \in T$,

$$\text{if } v \lesssim o \lesssim \iota, \text{ then } o = o \hat{\ominus} o.$$

Let $v \lesssim o \lesssim \iota$. Then we have $o = o \ominus o$ and so

$$\sup_{r \in T: r \leq o} r = \sup_{r \in T: r \leq o} \sup_{r \in T: r \leq o} r \ominus r.$$

Hence the claim holds. \square

Theorem 4.4. (Standard completeness) *Let Γ be a theory over $\mathbf{WA}_t\mathbf{BUL}$ and R be a formula. For $\mathbf{WA}_t\mathbf{BUL}$, the following are equivalent:*

1. $\Gamma \vdash_{\mathbf{WA}_t\mathbf{BUL}} R$.
2. For every standard $\mathbf{WA}_t\mathbf{BUL}$ -algebra and interpretation It , if $It(P) \geq \iota$ for all $P \in \Gamma$, then $It(R) \geq \iota$.

Proof. 1 \rightarrow 2: This follows from the definition.

2 \rightarrow 1: Let R be a formula such that $\Gamma \not\vdash_{\mathbf{WA}_t\mathbf{BUL}} R$, \mathbf{S} a linearly ordered $\mathbf{WA}_t\mathbf{BUL}$ -algebra, and It an interpretation in \mathbf{S} such that $It(P) \geq \iota$ for all $P \in \Gamma$ and $It(R) < \iota$. Let i' be the embedding of \mathbf{S} into the standard $\mathbf{WA}_t\mathbf{BUL}$ -algebra as in Proposition ???. Then, $i' \ominus It$ is an interpretation into the standard $\mathbf{WA}_t\mathbf{BUL}$ -algebra such that $i' \ominus It(P) \geq \iota$ and yet $i' \ominus It(R) < \iota$. \square

The proof of standard completeness for $\mathbf{WA}_t\mathbf{BUL}$ can be analogously applied to the other logics.

Theorem 4.5. *Let Γ be a theory over $\mathbf{WA}_t\mathbf{GBUL}$ and a finite theory over each of $\mathbf{WA}_t\mathbf{IBUL}$, $\mathbf{WA}_t\mathbf{CBUL}$ and $\mathbf{WA}_t\mathbf{CRL}$. For $\mathbf{WA}_t\mathbf{L} \in \{\mathbf{WA}_t\mathbf{IBUL}, \mathbf{WA}_t\mathbf{CBUL}, \mathbf{WA}_t\mathbf{GBUL}, \mathbf{WA}_t\mathbf{CRL}\}$, the following are equivalent:*

1. $\Gamma \vdash_{\mathbf{WA}_t\mathbf{L}} R$.
2. For every standard $\mathbf{WA}_t\mathbf{L}$ -algebra and interpretation It , if $It(P) \geq \iota$ for all $P \in \Gamma$, then $It(R) \geq \iota$.

Proof. First, the definition of \ominus for $\mathbf{WA}_t\mathbf{CBUL}$ is the same as in the proof of Proposition ???. Thus, for $\mathbf{WA}_t\mathbf{CBUL}$, we need to prove the conditions (GN^S) , (CP^S) and (RCAN_U^wS) . For a set T , Gödel negation \neg_G is defined as follows: for $(\sigma, o) \in T$,

$$\neg_G(\sigma, o) = \begin{cases} \bigwedge & \text{if } (\sigma, o) \neq \bigwedge \\ \bigvee & \text{otherwise} \end{cases}$$

For (GN^S) , we need to show: for $(\sigma, o) \in T$,

$$\min\{(\sigma, o), \neg_G(\sigma, o)\} = \bigwedge.$$

We consider the case $(\sigma, o) \neq \bigwedge$. From the definition, it is immediate that $\neg_G(\sigma, o) = \bigwedge$. Hence the claim holds.

For (CP^S) , we need to show: for $(\sigma, o), (\theta, l) \in T$,

$$\iota \lesssim ((\sigma, o) \rightarrow (\theta, l)) \rightarrow (\neg_G(\theta, l) \rightarrow \neg_G(\sigma, o)).$$

We instead show that

$$(\sigma, o) \rightarrow (\theta, l) \lesssim \neg_G(\theta, l) \rightarrow \neg_G(\sigma, o).$$

For $(\tau, o), (\sigma, l) \in T$, we may define $(\tau, o) \rightarrow (\sigma, l)$ as $\sup\{(\theta, m) \in T : (\tau, o) \ominus (\theta, m) \leq (\sigma, l)\}$. Let $(\sigma, o) = \bigwedge$. Then since $\neg_G(\sigma, o) = \bigvee$ and so $\neg_G(\theta, l) \rightarrow \neg_G(\sigma, o) = \bigvee$, the claim holds. Let $(\sigma, o) \neq \bigwedge$. If $(\theta, l) = \bigwedge$, then $(\sigma, o) \rightarrow (\theta, l) = \bigwedge$ and so the claim holds. Otherwise, $\neg_G(\sigma, o) = \neg_G(\theta, l) = \bigwedge$ and so $\neg_G(\theta, l) \rightarrow \neg_G(\sigma, o) = \bigvee$. Hence the claim holds.

For (RCAN_U^wS) , we need to show: for $(\sigma, o), (\theta, l) \in T$,

$$\iota \lesssim \min\{\neg_G\neg_G(\sigma, o), \iota\} \rightarrow (\min\{(\min\{(\sigma, o), v\} \rightarrow (\min\{(\sigma, o), v\} \ominus (\theta, l))), v\} \rightarrow (\theta, l)).$$

We instead show that

$$\min\{\neg_G\neg_G(\sigma, o), \iota\} \lesssim \min\{(\min\{(\sigma, o), v\} \rightarrow (\min\{(\sigma, o), v\} \ominus (\theta, l))), v\} \rightarrow (\theta, l).$$

Let $(\sigma, o) = \bigwedge$. Then $\neg_G\neg_G(\sigma, o) = \bigwedge$ and so $\min\{\neg_G\neg_G(\sigma, o), \iota\} = \bigwedge$. Hence the claim holds. Let $(\sigma, o) \neq \bigwedge$. Then we have $\neg_G\neg_G(\sigma, o) = \bigvee$ and so $\min\{\neg_G\neg_G(\sigma, o), \iota\} = \iota$. Thus we need to verify

$$(A) \quad \min\{(\min\{(\sigma, o), v\} \rightarrow (\min\{(\sigma, o), v\} \ominus (\theta, l))), v\} \lesssim (\theta, l).$$

We must consider the case $(\theta, l) \prec v$ since $\min\{(\min\{(\sigma, o), v\} \rightarrow (\min\{(\sigma, o), v\} \ominus (\theta, l))), v\} \lesssim v$. Note that $(\sigma_u \rightarrow (\sigma_u \star \theta))_u \leq \theta$. Thus one can assure that (A) holds.

For $\mathbf{WA}_t\mathbf{GBUL}$, we need to prove (RID_U^S) , i.e., $(\sigma, o) = (\sigma, o) \ominus (\sigma, o)$ for $(\sigma, o) \lesssim v$. Note that $\sigma \star \sigma = \sigma$. This ensures $(\sigma, o) = \min\{(\sigma, o), (\sigma, o)\} = (\sigma, o) \ominus (\sigma, o)$.

For $\mathbf{WA}_t\mathbf{IBUL}$, first take S as in the proof of Proposition ???. For each $\sigma \in S$, let σ^+ denote the successor of σ in case it exists, otherwise $\sigma^+ = \sigma$. Notice that, since the negation in S , defined as $\sim \sigma := \sigma \rightarrow f$, is involutive, one has that $\sigma = (\sim \theta)^+$ if and only if $\theta = (\sim \sigma)^+$; moreover, if $\sigma < \sigma^+$, then $(\sim (\sigma^+))^+ = \sim \sigma$. Let us take V below instead of the T , which is defined in the proof of Proposition ???. Let (V, \lesssim) be the linearly ordered set defined by

$$V = \{(\sigma, \sigma) : \sigma \in S\} \cup \{(\sigma, o) : \exists \sigma' \in S \text{ such that } \sigma = \sigma'^+ > \sigma', \text{ and } o \in \mathbf{Q} \cap (0, \sigma)\},$$

and \lesssim , which is the corresponding lexicographic order as above. Certainly (V, \lesssim) is a subset of the ordered set (T, \lesssim) defined in Proposition ??? having the same bounds and special elements ι, ρ and v . Note that V is closed under \ominus . Certainly, as in the proof of Proposition ???, one can verify the condition 1 (with V instead of T) in Proposition ???.

A new operation \odot on V , based on \ominus , is define as follows:

$$(\sigma, o) \odot (\theta, l) = \begin{cases} \max\{(\sigma + \theta - u, o + l - u), \bigwedge\} & \text{if } \sigma, \theta \leq u, \\ \min\{\partial, (\sigma, o) \ominus (\theta, l)\} & \text{if } \sigma > t \text{ or } \theta > t, \text{ and either} \\ & \sigma = (\sim \theta)^+ \text{ and } \frac{c}{d} + \frac{c'}{d'} \leq 1, \\ & \text{where } o = \sigma \frac{c}{d} \text{ and } l = \theta \frac{c'}{d'}, \text{ or} \\ & \sigma < (\sim \theta)^+; \\ (\sigma, o) \ominus (\theta, l) & \text{otherwise.} \end{cases}$$

We need to verify that the conditions (DNE^S) and (DNE_U^S) hold. For (DNE^S) , see Theorem 5 in [?]. Let $\neg(\sigma, o)$ be $(\sigma \rightarrow 0, o \rightarrow 0)$ for $(\sigma, o) \in T$. For (DNE_U^S) , we have to show: for $(\sigma, o) \in T$,

$$\iota \lesssim \min\{\neg\neg(\sigma, o), v\} \rightarrow (\sigma, o).$$

We instead show that

$$(B) \min\{\neg\neg(\sigma, o), v\} \lesssim (\sigma, o).$$

We need to consider the case $(\sigma, o) \lesssim v$. Let $\neg(\tau, o)$ be $(u - \tau, u - o)$ for $(\tau, o) \in V$ such that $(\tau, o) \lesssim v$. Then we have $\neg(\sigma, o) = (u - \sigma, u - o)$ and so $\neg\neg(\sigma, o) = (u - (u - \sigma), u - (u - o)) = (\sigma, o)$. Hence (B) holds.

For **WA_tCRL**, first note that the set V and operation \odot are the same as in **WA_tIBUL**. We need to verify that the condition (FP^S) holds. For (FP^S), we have to show:

$$\iota = \rho.$$

This is immediate since $t = f$ and so $(t, t) = (f, f)$.

The proof of the remaining for standard completeness of **WA_tL** is almost the same as in **WA_tBUL**. \square

Let us call a linearly ordered algebra a *chain*. We note that every **WA_tBL**-chain (**WA_tL**-chain, **WA_tΠ**-chain and **WA_tG**-chain resp) contains a **BL**-chain (**L**-chain, **P**-chain and **G**-chain resp) (see Proposition 5 in [?]). Analogously, we show that every **WA_tBUL**-chain (**WA_tIBUL**-chain, **WA_tCBUL**-chain and **WA_tGBUL**-chain resp) contains a **BL**-chain (**L**-chain, **P**-chain and **G**-chain resp). Let S be the universe and $S_u := \{o : \perp \leq o \leq u\}$. Define $o \rightarrow_u l := (o \rightarrow l) \wedge u$.

Proposition 4.6. 1. For every **WA_tBUL**-chain, $(S_u, \wedge, \vee, \star, \rightarrow_u, \perp, u)$ is a **BL**-chain.

2. For every **WA_tIBUL**-chain, $(S_u, \wedge, \vee, \star, \rightarrow_u, \perp, u)$ is a **L**-chain.

3. For every **WA_tCBUL**-chain, $(S_u, \wedge, \vee, \star, \rightarrow_u, \perp, u)$ is a **P**-chain.

4. For every **WA_tGBUL**-chain, $(S_u, \wedge, \vee, \star, \rightarrow_u, \perp, u)$ is a **G**-chain.

Proof. 1. If $o, l \in S_u$, then it is clear that $o \wedge l, o \vee l, o \star l, o \rightarrow_u l \in S_u$. Moreover, $(S_u, \wedge, \vee, \perp, u)$ is a bounded lattice and (S_u, \star, u) is a commutative monoid. It is routine to verify prelinearity and residuation properties. We verify divisibility property. One has to show: for all $o, l \in S_u$,

$$u = (o \rightarrow_u l) \vee (l \rightarrow_u (o \star (o \rightarrow_u l))).$$

Let $o \leq l$. Then since $u = o \rightarrow_u l$ by (residuation), we are done. Let $o > l$. Then, since $l < o \leq u$, we have $t \leq l \rightarrow (o \star (o \rightarrow l))$ by (RDIV_t^{wS}) and so $u \leq l \rightarrow (o \star (o \rightarrow l))$. Since $l < o = o \star u$ by (U-RI^S) and so $o \rightarrow l < u$, we have $o \rightarrow l = o \rightarrow_u l$ and so $u \leq l \rightarrow_u (o \star (o \rightarrow_u l))$. Hence $u = l \rightarrow_u (o \star (o \rightarrow_u l))$.

2. By 1, we need to verify that the negation is involutive. Let us define $\neg_u o = o \rightarrow_u \perp$. Then, for all $o, l \in S_u$, we can define $o \star l := \neg_u(o \rightarrow_u \neg_u l)$ and $o \rightarrow_u l := \neg_u(o \star \neg_u l)$. Since $\neg_u(o \rightarrow_u \perp) = o \star u = o$, we have $\neg_u \neg_u o = o$.

3. Let \neg_G be the \neg_u satisfying (GN^S). We need to verify cancellation property, i.e., for all $o, l \in S_u$,

$$u = \neg_G \neg_G o \rightarrow_u ((o \rightarrow_u (o \star l)) \rightarrow_u l).$$

Suppose $o = \perp$. We have $\neg_G \neg_G o = \perp$ and so the claim holds. Otherwise, $\neg_G \neg_G o = u$. Then we have to verify

$$u = (o \rightarrow_u (o \star l)) \rightarrow_u l.$$

Let $o \leq o \star l$. By (RCAN_U^{wS}), we obtain $u \leq ((o \rightarrow (o \star l)) \wedge u) \rightarrow l$, and so $u = (o \rightarrow_u (o \star l)) \rightarrow_u l$. Otherwise, we get $t \leq ((o \rightarrow (o \star l)) \wedge u) \rightarrow l$ by (RCAN_U^{wS}). Since $o \star l < o = o \star u$ by (U-RI^S), we further obtain $o \rightarrow (o \star l) = o \rightarrow_u (o \star l)$ and so $u = (o \rightarrow_u (o \star l)) \rightarrow_u l$. Therefore the claim holds.

4. We further need to verify (RID_U^S). Since $o \star o = o \leq u$ for all $o, l \in S_u$, we have $o \star o = (o \star o) \wedge u = o \wedge u$. Hence \star satisfies (RID_U^S). \square

We finally verify that each standard **WA_tL**-algebra is given over a restricted v -identical ι_v -idempotent $[0, t]$ -continuous wa_t -uninorm.

Proposition 4.7. Let $\mathcal{S} = ([0, 1], 1, 0, \iota, \rho, v, \min, \max, \ominus, \rightarrow)$ be a **WA_tL**-algebra. The operation \ominus is $[0, \iota]$ -continuous, restricted v -identical and ι_v -idempotent.

Proof. Let \mathcal{S} be a **WA_tL**-algebra. For $[0, \iota]$ -continuity of \ominus , we first notice that a wa_t -uninorm \ominus is residuated if and only if it is conjunctive and left-continuous (see [?]). Hence we need to verify that \ominus is right-continuous on $[0, \iota]$. Suppose that $o, l \in [0, \iota]$ and $(o_i)_{0 \leq i}$ is a non-increasing sequence in $[0, \iota]$ so that $o = \inf_i o_i$. If $m = \inf_i (o_i \ominus l)$, then $o \ominus l \leq m$. Thus, one has to verify $o \ominus l \geq m$. Notice that the monoid operation of a standard **BL**-algebra (**P**-algebra, **L**-algebra, **G**-algebra resp) forms a continuous **t**-norm. Hence \ominus is continuous on $[0, u]$ by Proposition ???. Moreover, since $\ominus = \min$ on $[v, \iota]$ by (ID_U^{tS}) and \min is continuous, \ominus is continuous on $[v, \iota]$. Therefore, \ominus is continuous on $[0, \iota]$.

For the restricted v -identity of the operation \ominus , let $o \in [0, 1]$ be such that $o \leq v$. Then $v \ominus o = o$ by (U-RI^S). Hence it is restricted v -identical.

For the ι_v -idempotence of the operation \ominus , let $o \in [0, 1]$ be such that $v \leq o \leq \iota$. Then $o \ominus o = o$ by (ID_U^{tS}). Hence it is ι_v -idempotent. \square

Theorem 4.8. 1. (*Finite standard completeness*) Let Γ be a finite theory over WA_tL ($\in Ls$) and R a formula. $\Gamma \vdash_{WA_tL} R$ if and only if for each standard $[0, \iota]$ -continuous ι_v -idempotent restricted v -identical \mathbf{WA}_tL -algebra and interpretation It , if $It(P) \geq \iota$ for each $P \in \Gamma$, then $It(R) \geq \iota$.

2. (*Standard completeness*) Let Γ be a theory over $WA_tL \in \{\mathbf{WA}_tBUL, \mathbf{WA}_tGBUL\}$ and R a formula. $\Gamma \vdash_{WA_tL} R$ if and only if for each standard $[0, \iota]$ -continuous ι_v -idempotent restricted v -identical \mathbf{WA}_tL -algebra and interpretation It , if $It(P) \geq \iota$ for each $P \in \Gamma$, then $It(R) \geq \iota$.

Proof. 1 and 2 follow from Propositions ?? and ??, and Theorems ?? and ??. \square

5 Concluding remarks

Micanorm-based logics with a weak form of associativity, denoted by basic wa_t -uninorm logics, were introduced as a weak associative generalization of basic uninorm logics introduced by Gabbay and Metcalfe [?] and it was verified that those logics are algebraically complete. After related uninorms with the weak associativity instead of associativity were introduced as $[0, t]$ -continuous wa_t -uninorms, it was proved that the logics are standard complete on these wa_t -uninorms.

Notice that Yang [?] introduced such a generalization of continuous t -norm-based logics and left the work in this paper as an open problem. This is the answer to the problem. Furthermore, it will provide an insight for algebraic applications based on weak associative logics.

We may think of other weak associative generalizations of basic uninorm logics. For instance, t -associative and strong t -associative generalizations can be introduced as in [?]. Moreover, a weak u -associative generalization can be dealt with in place of the weak t -associative generalization addressed in this paper. (Note that the basic uninorm logics were introduced as logics based on $[0, 1]$ -continuous uninorms, which have the element u , see [?].) To study such systems and related semantics is a future work. Moreover, as a problem it remains to investigate applications such as weak t -associative compensation behavior and reinforcement in wa_t -uninorms.

Acknowledgment

The author appreciates the anonymous referees for their helpful comments and suggestions to improve the manuscript.

References

- [1] A. A. Adrian, *Power-associative rings*, Transactions of the American Mathematical Society, **64** (1948), 552-593.
- [2] M. Christos, C. Tsitouras, *Enumeration of Rosenberg type hypercompositional structures defined by binary relations*, Italian Journal of Pure and Applied Mathematics, **28** (2011), 43-54. <https://doi.org/10.1016/j.ejcm.2012.03.032>
- [3] J. Chvalina, Š. Hošková-Mayerová, *On certain proximities and preorderings on the transposition hypergroups of linear first-order partial differential operators*, Analele științifice ale Universitatii Ovidius Constanta, **22** (2014), 85-103. <https://doi.org/10.2478/auom-2014-0008>
- [4] P. Cintula, R. Horčík, C. Noguera, *Non-associative substructural logics and their semilinear extensions: axiomatization and completeness properties*, Review of Symbolic Logic, **6** (2013), 394-423. <https://doi.org/10.1017/S1755020300000000>
- [5] P. Cintula, R. Horčík, C. Noguera, *The quest for the basic fuzzy logic*, in Petr Hájek on Mathematical Fuzzy Logic, F. Montagna, (ed), Springer, Dordrecht (2015), 245-290. https://doi.org/10.1007/978-3-319-06233-4_12
- [6] P. Cintula, C. Noguera, *A general framework for mathematical fuzzy logic*, in Handbook of Mathematical Fuzzy Logic, Vol. 1, P. Cintula, R. Horčík, C. Noguera (eds.), College Publications, London, (2011), 103-207.

- [7] F. Esteva, L. Godo, *Monoidal t -norm based logic: Towards a logic for left-continuous t -norms*, Fuzzy Sets and Systems, **124** (2001), 271-288. [https://doi.org/10.1016/S0165-0114\(01\)00098-7](https://doi.org/10.1016/S0165-0114(01)00098-7)
- [8] F. Esteva, L. Gispert, L. Godo, F. Montagna, *On the standard and rational completeness of some axiomatic extensions of the monoidal t -norm logic*, Studia Logica, **71** (2002), 393-420. <https://doi.org/10.1023/A:1016548805869>
- [9] E. Fried, G. Grätzer, *A nonassociative extensions of the class of distributive lattices*, Pacific Journal of Mathematics, **49** (1973), 59-78. <https://projecteuclid.org/journals/pacific-journal-of-mathematics/volume-49/issue-1/A-nonassociative-extension-of-the-class-of-distributive-lattices/pjm/1102945268.full>
- [10] D. Gabbay, G. Metcalfe, *Fuzzy Logics based on $[0, 1)$ -continuous uninorms*, Archive for Mathematical Logic, **46** (2007), 425-449. <https://doi.org/10.1007/s00153-007-0047-1>
- [11] M. Ghadiri, B. Davvaz, *Direct system and direct limit of Hv-modules*, Iranian Journal of Science and Technology, Transaction A, **28** (2004), 267-275.
- [12] G. Grätzer, A. Kisielewicz, B. Wolk, *An equational basis in four variables for the three-element tournament*, Colloquium Mathematicum, **18** (1992), 41-44. <https://doi.org/10.4064/cm-63-1-41-44>
- [13] P. Hájek, *Metamathematics of fuzzy logic*, Kluwer, Amsterdam, 1998. <https://doi.org/10.1007/978-94-011-5300-3>
- [14] S. Jenei, F. Montagna, *A proof of standard completeness for Esteva and Godo's logic MTL*, Studia Logica, **70** (2002), 183-192. <https://doi.org/10.1023/A:1015122331293>
- [15] T. Kowalski, *Weakly associative relation algebras hold the key to the universe*, Bulletin of the Section of Logic, **36** (2007), 145-157. <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=4bc2455a421b9fbc2ff63aaa49113ba907f90c45>
- [16] K. Kunnen, *Quasigroups, loops, and associative laws*, Journal of Algebra, **185** (1996), 194-204. <https://doi.org/10.1006/jabr.1996.0321>
- [17] A. Kurucz, *Weakly associative relation algebras with projections*, Mathematical Logic Quarterly, **29** (2008), 1-16. <https://doi.org/10.1002/malq.200710074>
- [18] N. Lygeros, T. Vougiouklis, *The LV-hyperstructures*, Ratio Mathematica, **25** (2013), 59-66. https://lygeros.org/wp-content/uploads/2019/12/14279_RM_25_5.pdf
- [19] G. Metcalfe, F. Montagna, *Substructural fuzzy logics*, Journal of Symbolic Logic, **72** (2007), 834-864. <https://www.jstor.org/stable/27588573>
- [20] G. Metcalfe, N. Olivetti, D. Gabbay, *Proof theory for fuzzy logics*, Springer Series in Applied Logic, vol. **36**, 2009. <https://doi.org/10.1007/978-1-4020-9409-5>
- [21] B. O. Onasanya, T. S. Atamewoue, S. Hoskova-Mayerova, *Certain fuzzy hyperstructures from a fuzzy set*, Journal of Intelligent and Fuzzy Systems, **39** (2020), 2775-2782. [10.3233/JIFS-191054](https://doi.org/10.3233/JIFS-191054)
- [22] J. M. Pallo, *Weak associativity and restricted rotation*, Information Processing Letters, **109** (2009), 514-517. <https://doi.org/10.1016/j.ipl.2009.01.014>
- [23] G. Restall, G. *An introduction to substructural logics*, New York, Routledge, 2000.
- [24] D. Šalounová, *Spectra of weakly associative lattice rings*, Acta Universitatis Palackianae Olomucensis Facultas Rerum Naturalium Mathematica, **40** (2001), 215-224.
- [25] R. D. Schafer, *An introduction to non-associative algebras*, Dover, 1996. <https://www.gutenberg.org/files/25156/25156-pdf.pdf>
- [26] V. Stebletsova, *Weakly associative relation algebras with polyadic composition operations*, Studia Logica, **66** (2000), 297-323. <https://doi.org/10.1023/A:1005204532371>

- [27] O. Susumu, *Introduction to octonion and other non-associative algebras in physics (Montroll Memorial Lecture Series in Mathematical Physics)*, Cambridge, Cambridge Univ. Press, 1995. <https://doi.org/10.1017/CBO9780511524479>
- [28] T. Vougiouklis, *The e-hyperstructures*, Journal of Mahani Mathematical Research Center, **1** (2012), 13-28. <https://doi.org/10.22103/JMMRC.2012.345>
- [29] T. Vougiouklis, *Enlarged fundamentally very thin Hv-structures*, Journal of Algebraic Structures and Their Applications, **1** (2014), 11-20. https://as.yazd.ac.ir/article_410_a84741ff38a501826228c29a144726fd.pdf
- [30] T. Vougiouklis, S. Spartalis, M. Kessoglides, *Weak hyperstructures on small sets*, Ratio Mathematica, **12** (1997), 90-96.
- [31] S. Wang, *Uninorm logic with the n-potency axiom*, Fuzzy Sets and Systems, **205** (2012), 116-126. <https://doi.org/10.1016/j.fss.2012.04.017>
- [32] S. Wang, *Involutive uninorm logic with the n-potency axiom*, Fuzzy Sets and Systems, **218** (2013), 1-23. <https://doi.org/10.1016/j.fss.2012.09.009>
- [33] E. Yang, *Weakening-free, non-associative fuzzy logics: Micanorm-based logics*, Fuzzy Sets and Systems, **276** (2015), 43-58. <https://doi.org/10.1016/j.fss.2014.11.020>
- [34] E. Yang, *Basic substructural core fuzzy logics and their extensions: Mianorm-based logics*, Fuzzy Sets and Systems, **301** (2016), 1-18. <https://doi.org/10.1016/j.fss.2015.09.007>
- [35] E. Yang, *Weakly associative fuzzy logics*, Korean Journal of Logic, **19** (2016), 437-461.
- [36] E. Yang, *Involutive basic substructural core fuzzy logics: Involutive mianorm-based logics*, Fuzzy Sets and Systems, **320** (2017), 1-16. <https://doi.org/10.1016/j.fss.2017.03.013>
- [37] E. Yang, *A non-associative generalization of continuous t-norm-based logics*, Journal of Intelligent and Fuzzy Systems, **33** (2017), 3743-3752. <https://doi.org/10.3233/JIFS-17623>
- [38] E. Yang, *Involutive micanorm logics with n-potency axiom*, Korean Journal of Logic, **20** (2017), 273-292. <https://logicalkorea.com/attach/paper/20-2-ESYang.pdf?ckattempt=1>
- [39] E. Yang, *Mianorm-based logics with n-contraction and n-mingle axioms*, Journal of Intelligent and Fuzzy Systems, **37** (2019), 7895-7907. <https://doi.org/10.3233/JIFS-190150>
- [40] E. Yang, *Involutive extensions of fixpointed mianorm-based logics*, Korean Journal of Logic, **24** (2021), 31-52.
- [41] E. Yang, *Micanorm-based logics with fixed-point*, Korean Journal of Logic, **24** (2021), 281-297.
- [42] E. Yang, *Fixed-pointed involutive micanorm-based logics*, Korean Journal of Logic, **25** (2022), 121-137. https://logicalkorea.com/attach/paper/25-2_01_YangES.pdf
- [43] E. Yang, *Semilinear logics with knotted axioms*, Iranian Journal of Fuzzy Systems, **19** (2022), 17-30. <https://doi.org/10.22111/IJFS.2022.6785>